



## Complete Summary

---

### TITLE

Heart failure in adults: percentage of adult patients with a primary diagnosis of heart failure who have used tobacco anytime during the year prior to hospital arrival and who are given smoking cessation advice or counseling during the hospital stay or at discharge.

### SOURCE(S)

Institute for Clinical Systems Improvement (ICSI). Heart failure in adults. Bloomington (MN): Institute for Clinical Systems Improvement (ICSI); 2007 Aug. 119 p. [217 references]

## Measure Domain

### PRIMARY MEASURE DOMAIN

Process

The validity of measures depends on how they are built. By examining the key building blocks of a measure, you can assess its validity for your purpose. For more information, visit the [Measure Validity](#) page.

### SECONDARY MEASURE DOMAIN

Does not apply to this measure

## Brief Abstract

### DESCRIPTION

This measure is used to assess the percentage of adult patients with a primary diagnosis of heart failure who have used tobacco anytime during the year prior to hospital arrival and who are given smoking cessation advice or counseling during the hospital stay or at discharge.

### RATIONALE

The priority aim addressed by this measure is to improve care of adult heart failure patients by assuring comprehensive patient education and follow-up care.

### PRIMARY CLINICAL COMPONENT

Heart failure; smoking cessation advice or counseling

#### **DENOMINATOR DESCRIPTION**

Number of adult patients with a primary diagnosis of heart failure who have used tobacco anytime during the year prior to hospital arrival (see the related "Denominator Inclusions/Exclusions" field in the Complete Summary)

#### **NUMERATOR DESCRIPTION**

Number of adult patients with a primary diagnosis of heart failure who have used tobacco anytime during the year prior to hospital arrival and who are given smoking cessation advice or counseling during the hospital stay or at discharge

### **Evidence Supporting the Measure**

#### **EVIDENCE SUPPORTING THE CRITERION OF QUALITY**

- A clinical practice guideline or other peer-reviewed synthesis of the clinical evidence

#### **NATIONAL GUIDELINE CLEARINGHOUSE LINK**

- [Heart failure in adults.](#)

### **Evidence Supporting Need for the Measure**

#### **NEED FOR THE MEASURE**

Unspecified

### **State of Use of the Measure**

#### **STATE OF USE**

Current routine use

#### **CURRENT USE**

Internal quality improvement

### **Application of Measure in its Current Use**

#### **CARE SETTING**

Hospitals

#### **PROFESSIONALS RESPONSIBLE FOR HEALTH CARE**

Physicians

#### **LOWEST LEVEL OF HEALTH CARE DELIVERY ADDRESSED**

Group Clinical Practices

#### **TARGET POPULATION AGE**

Age greater than or equal to 18 years

#### **TARGET POPULATION GENDER**

Either male or female

#### **STRATIFICATION BY VULNERABLE POPULATIONS**

Unspecified

### **Characteristics of the Primary Clinical Component**

#### **INCIDENCE/PREVALENCE**

Heart failure is a major health problem in the United States, and the incidence of the disease is projected to increase. It was the most frequent diagnosis of Medicare patients discharged from the hospital in 2001. There are an estimated 5 million individuals currently diagnosed with heart failure, 550,000 new cases diagnosed annually, and 1 million people hospitalized annually (including readmission rates of 30 to 60%).

#### **EVIDENCE FOR INCIDENCE/PREVALENCE**

Institute for Clinical Systems Improvement (ICSI). Heart failure in adults. Bloomington (MN): Institute for Clinical Systems Improvement (ICSI); 2007 Aug. 119 p. [217 references]

#### **ASSOCIATION WITH VULNERABLE POPULATIONS**

Unspecified

#### **BURDEN OF ILLNESS**

There are over 260,000 annual deaths from heart failure. From 1979 to 2000 heart failure death rates have increased 148% and hospital discharges have increased 165%. The prognosis of patients with a new diagnosis of heart failure is poor. Senni et al. (1998) noted survival to be 86% at 3 months, 76% at one year, and only 35% at 5 years.

#### **EVIDENCE FOR BURDEN OF ILLNESS**

Institute for Clinical Systems Improvement (ICSI). Heart failure in adults. Bloomington (MN): Institute for Clinical Systems Improvement (ICSI); 2007 Aug. 119 p. [217 references]

Senni M, Tribouilloy CM, Rodeheffer RJ, Jacobsen SJ, Evans JM, Bailey KR, Redfield MM. Congestive heart failure in the community: a study of all incident cases in Olmsted County, Minnesota, in 1991. *Circulation* 1998 Nov 24;98(21):2282-9. [PubMed](#)

## **UTILIZATION**

See the "Incidence/Prevalence" field.

## **COSTS**

Unspecified

## **Institute of Medicine National Healthcare Quality Report Categories**

### **IOM CARE NEED**

Living with Illness

### **IOM DOMAIN**

Effectiveness  
Patient-centeredness

## **Data Collection for the Measure**

### **CASE FINDING**

Users of care only

### **DESCRIPTION OF CASE FINDING**

Adult patients with a primary diagnosis of heart failure who have used tobacco anytime during the year prior to hospital arrival and who are discharged alive.

The time of assessment is at hospital discharge. Monthly data will be submitted quarterly.

The minimum sample size is 20 patients per month.

### **DENOMINATOR SAMPLING FRAME**

Patients associated with provider

### **DENOMINATOR INCLUSIONS/EXCLUSIONS**

**Inclusions**

Number of adult patients with a primary diagnosis of heart failure\* who have used tobacco anytime during the year prior to hospital arrival\*\*.

\*International Classification of Diseases, Ninth Revision (ICD-9) codes: 402.01, 402.11, 402.91, 404.01, 404.03, 404.11, 404.13, 404.91, 404.93, 428.0, 428.1, 428.20, 428.21, 428.22, 428.23, 428.30, 428.31, 428.32, 428.33, 428.40, 428.41, 428.42, 428.43, 428.9. (Refer to the table, "Descriptions of ICD-9 Codes," in the original measure documentation for code descriptions.)

\*\*Tobacco user is defined as someone who has smoked tobacco any time during the year prior to hospitalization.

**Exclusions**

- Patients who are less than 18 years of age
- Patients who died prior to discharge
- Patients who were transferred to another hospital
- Patients who left against medical advice
- Patients discharged to hospice

**RELATIONSHIP OF DENOMINATOR TO NUMERATOR**

All cases in the denominator are equally eligible to appear in the numerator

**DENOMINATOR (INDEX) EVENT**

Clinical Condition  
Institutionalization

**DENOMINATOR TIME WINDOW**

Time window is a single point in time

**NUMERATOR INCLUSIONS/EXCLUSIONS****Inclusions**

Number of adult patients with a primary diagnosis of heart failure who have used tobacco anytime during the year prior to hospital arrival and who are given smoking cessation advice or counseling during the hospital stay or at discharge

**Exclusions**

Unspecified

**MEASURE RESULTS UNDER CONTROL OF HEALTH CARE PROFESSIONALS, ORGANIZATIONS AND/OR POLICYMAKERS**

The measure results are somewhat or substantially under the control of the health care professionals, organizations and/or policymakers to whom the measure applies.

**NUMERATOR TIME WINDOW**

Institutionalization

**DATA SOURCE**

Administrative data  
Medical record

**LEVEL OF DETERMINATION OF QUALITY**

Individual Case

**PRE-EXISTING INSTRUMENT USED**

Unspecified

**Computation of the Measure**

**SCORING**

Rate

**INTERPRETATION OF SCORE**

Better quality is associated with a higher score

**ALLOWANCE FOR PATIENT FACTORS**

Unspecified

**STANDARD OF COMPARISON**

Internal time comparison

**Evaluation of Measure Properties**

**EXTENT OF MEASURE TESTING**

Unspecified

**Identifying Information**

**ORIGINAL TITLE**

Percentage of adult patients with a primary diagnosis of heart failure who have used tobacco anytime during the year prior to hospital arrival and who are given smoking cessation advice or counseling during the hospital stay or at discharge.

**MEASURE COLLECTION**

## [Heart Failure in Adults Measures](#)

### **DEVELOPER**

Institute for Clinical Systems Improvement

### **ADAPTATION**

Measure was not adapted from another source.

### **RELEASE DATE**

2006 Aug

### **REVISION DATE**

2007 Aug

### **MEASURE STATUS**

This is the current release of the measure.

This measure updates a previous version: Institute for Clinical Systems Improvement (ICSI). Heart failure in adults. Bloomington (MN): Institute for Clinical Systems Improvement (ICSI); 2006 Aug. 116 p.

### **SOURCE(S)**

Institute for Clinical Systems Improvement (ICSI). Heart failure in adults. Bloomington (MN): Institute for Clinical Systems Improvement (ICSI); 2007 Aug. 119 p. [217 references]

### **MEASURE AVAILABILITY**

The individual measure, "Percentage of adult patients with a primary diagnosis of heart failure who have used tobacco anytime during the year prior to hospital arrival and who are given smoking cessation advice or counseling during the hospital stay or at discharge," is published in "Health Care Guideline: Heart Failure in Adults." This document is available from the [Institute for Clinical Systems Improvement \(ICSI\) Web site](#).

For more information, contact ICSI at, 8009 34th Avenue South, Suite 1200, Bloomington, MN 55425; phone: 952-814-7060; fax: 952-858-9675; Web site: [www.icsi.org](http://www.icsi.org); e-mail: [icsi.info@icsi.org](mailto:icsi.info@icsi.org).

### **NQMC STATUS**

This NQMC summary was completed by ECRI on November 24, 2006. This NQMC summary was updated by ECRI Institute on December 16, 2007.

## **COPYRIGHT STATEMENT**

This NQMC summary (abstracted Institute for Clinical Systems Improvement [ICSI] Measure) is based on the original measure, which is subject to the measure developer's copyright restrictions.

The abstracted ICSI Measures contained in this Web site may be downloaded by any individual or organization. If the abstracted ICSI Measures are downloaded by an individual, the individual may not distribute copies to third parties.

If the abstracted ICSI Measures are downloaded by an organization, copies may be distributed to the organization's employees but may not be distributed outside of the organization without the prior written consent of the Institute for Clinical Systems Improvement, Inc.

All other copyright rights in the abstracted ICSI Measures are reserved by the Institute for Clinical Systems Improvement, Inc. The Institute for Clinical Systems Improvement, Inc. assumes no liability for any adaptations or revisions or modifications made to the abstracts of the ICSI Measures.

### **Disclaimer**

## **NQMC DISCLAIMER**

The National Quality Measures Clearinghouse™ (NQMC) does not develop, produce, approve, or endorse the measures represented on this site.

All measures summarized by NQMC and hosted on our site are produced under the auspices of medical specialty societies, relevant professional associations, public and private organizations, other government agencies, health care organizations or plans, individuals, and similar entities.

Measures represented on the NQMC Web site are submitted by measure developers, and are screened solely to determine that they meet the NQMC Inclusion Criteria which may be found at <http://www.qualitymeasures.ahrq.gov/about/inclusion.aspx>.

NQMC, AHRQ, and its contractor ECRI Institute make no warranties concerning the content or its reliability and/or validity of the quality measures and related materials represented on this site. The inclusion or hosting of measures in NQMC may not be used for advertising or commercial endorsement purposes.

Readers with questions regarding measure content are directed to contact the measure developer.

Date Modified: 11/3/2008

